

Ramsey-like theorems for the Schreier barrier

Quentin Le Houérou
Université Paris-Est-Créteil-Val-de-Marne

CCR2025 $\oplus$ JC

June 19, 2025

Joint work with Lorenzo Carlucci, Oriola Gjetaj and Ludovic
Levy Patey

Introduction

Theorems as problems

Many theorems (of ordinary mathematics) can be seen as problems involving instances and solutions:

- For every field, there exists an algebraic closure.
- For infinite binary tree, there exists an infinite branch (Weak König's lemma)
- For every bounded sequence in $\mathbb{R}^n$, there exists a converging subsequence (Bolzano-Weierstrass theorem).

The proof of these theorems often give a "procedure" to build the solution given the instance.

We can then wonder about the effectiveness of such constructions.

Theorems as problems

Many theorems (of ordinary mathematics) can be seen as problems involving instances and solutions:

- For every field, there exists an algebraic closure.
- For infinite binary tree, there exists an infinite branch (Weak König's lemma)
- For every bounded sequence in $\mathbb{R}^n$, there exists a converging subsequence (Bolzano-Weierstrass theorem).

The proof of these theorems often give a "procedure" to build the solution given the instance.

We can then wonder about the effectiveness of such constructions.

Computational content of theorems

Given $X \subseteq \mathbb{N}$ an instance of a problem P , can we compute a solution $Y \subseteq \mathbb{N}$ from X ?

In a lot of cases the answer will be no:

- There exists computable consistent theories that have no computable models (e.g. Tennenbaum's theorem)
- There exists computable commutative rings with no computable maximal ideals.

Such theorems will therefore not be provable in constructive mathematics.

But we can push the computational analysis further and obtain more precise proof-theoretic results.

Computational content of theorems

Given $X \subseteq \mathbb{N}$ an instance of a problem P , can we compute a solution $Y \subseteq \mathbb{N}$ from X ?

In a lot of cases the answer will be no:

- There exists computable consistent theories that have no computable models (e.g. Tennenbaum's theorem)
- There exists computable commutative rings with no computable maximal ideals.

Such theorems will therefore not be provable in constructive mathematics.

But we can push the computational analysis further and obtain more precise proof-theoretic results.

Computational content of theorems

Given $X \subseteq \mathbb{N}$ an instance of a problem P , can we compute a solution $Y \subseteq \mathbb{N}$ from X ?

In a lot of cases the answer will be no:

- There exists computable consistent theories that have no computable models (e.g. Tennenbaum's theorem)
- There exists computable commutative rings with no computable maximal ideals.

Such theorems will therefore not be provable in constructive mathematics.

But we can push the computational analysis further and obtain more precise proof-theoretic results.

Reverse Mathematics

To prove that an axiom was needed for a theorem, we can try to prove the axiom from the theorem:

- Modulo ZF, Zorn's lemma implies the axiom of choice.
- Modulo absolute geometry, the Pythagorean theorem implies the Parallel postulate.

Reverse mathematics does the same, but modulo constructive mathematics.

We will use the formalism of second-order arithmetic.

All the models considered will be ω -models, i.e. models $(\mathbb{N}, S, +, \times, 0)$ with $(\mathbb{N}, +, \times, 0)$ the standard integers, and $S \subseteq 2^{\mathbb{N}}$ the second-order part.


Base theory (corresponding to computable mathematics) RCA_0 :

- Robinson's arithmetic Q
- Δ_1^0 -comprehension (The computable sets exists)
- Σ_1^0 -induction (Every set of finite cardinality is bounded)

The computable sets form a minimal ω -model of RCA_0 .

The “Big Five”

Modulo RCA_0 , most theorems of ordinary mathematics are equivalent to one the following theories (from weakest to strongest):



Definition (ACA₀)

ACA₀ is RCA₀ plus the comprehension axiom for every arithmetical formula.

This is equivalent to comprehension for Σ_1^0 formula, or to the existence of the Turing jump of any set.

The arithmetical sets form a minimal model ω -model of ACA₀.

For a problem P , if there exists an ω -model of $\text{RCA}_0 + P$ not containing the halting set K , then $\text{RCA}_0 + P \not\models \text{ACA}_0$.

Definition (ACA₀)

ACA₀ is RCA₀ plus the comprehension axiom for every arithmetical formula.

This is equivalent to comprehension for Σ_1^0 formula, or to the existence of the Turing jump of any set.

The arithmetical sets form a minimal model ω -model of ACA₀.

For a problem P , if there exists an ω -model of RCA₀ + P not containing the halting set K , then RCA₀ + $P \not\vdash$ ACA₀.

Cone avoidance

Definition (Cone avoidance)

A problem P admits *cone avoidance* if for every set Z , every non- Z -computable set C , every Z -computable instance X of P has a solution Y that does not compute C .

Proposition

*If P admits cone avoidance, then for every non computable set C , there exists a model of $\text{RCA}_0 + P$ that does not contains C .
Hence $\text{RCA}_0 + P \not\models \text{ACA}_0$.*


Strong cone avoidance

Definition (*strong* Cone avoidance)

A problem P admits *strong cone avoidance* if for every set Z , every non- Z -computable set C , every ~~Z -computable~~ instance X of P has a solution Y that does not compute C .


Cone avoidance witness the computable weakness of a problem P .
Strong cone avoidance witness its combinatorial weakness.

The infinite Ramsey's theorem for pairs and two colors escape the structure phenomenon of Ramsey's theory.




Ramsey Theory

Finite Ramsey's theorem



For every 2-coloring of the
edges of K_6

Finite Ramsey's theorem



There exists some
monochromatic copy of K_3

Infinite Ramsey's theorem

Let $[X]^n$ be the set of all subsets of X of cardinality n .

Definition (Ramsey's theorem)

RT_k^n is the statement: "For every coloring $f : [\mathbb{N}]^n \rightarrow k$, there is an infinite set $H \subseteq \mathbb{N}$ such that $|f([H]^n)| = 1$ ".

RT_k^1 is the infinite pigeonhole principle. It is computably true (albeit not by a uniform procedure, we can't computably guess which color to choose) and $\text{RCA}_0 \vdash \text{RT}_k^1$ for every $k \in \mathbb{N}$.

Facts

- RT_2^1 admits strong cone avoidance, i.e., for every set A and every non computable set Z , there exists some subset $H \subseteq A$ or $H \subseteq \bar{A}$ such that $H \not\leq_T Z$. (Dzhafarov/Jockusch)
- There exists a $\emptyset'$ -computable instance of RT_2^2 , every solution of which computes $\emptyset'$ (Hence RT_2^2 does not admits strong cone avoidance)
- RT_2^2 admits cone avoidance. (Dzhafarov/Jockusch)
- For every $n \geq 3$, there exists a computable instance of RT_2^n , every solution of which computes $\emptyset^{(n-2)}$. Hence $RCA_0 + RT_2^n \vdash ACA_0$ (Hirschfeldt/ Jockusch).

Ramsey-like theorems

What about weakenings of Ramsey's theorems ?

Definition (Thin set theorem)


Let TS^n be the statement: “for every coloring $f : [\mathbb{N}]^n \rightarrow \mathbb{N}$, there exists some infinite set $H \subseteq \mathbb{N}$ such that $f([H]^n) \neq \mathbb{N}$.”

Definition (Free set theorem)

Let FS^n be the statement: “for every coloring $f : [\mathbb{N}]^n \rightarrow \mathbb{N}$, there exists some infinite set $H \subseteq \mathbb{N}$, such that $f(s) \notin H \setminus s$ for every $s \in [H]^n$.”

Definition (Rainbow Ramsey theorem)

Let RRT_k^n be the statement: “for every coloring $f : [\mathbb{N}]^n \rightarrow \mathbb{N}$ such that $|f^{-1}(\ell)| \leq k$ for every $\ell \in \mathbb{N}$, there exists some infinite $H \subseteq \mathbb{N}$ such that f is injective on $[H]^n$.”



Theorem (Wang)

For every n, k , TS^n , FS^n and RRT_k^n admits the strong cone avoidance property.

Schreier barrier

Wrong generalizations

Let $[H]^{<\mathbb{N}}$ be the set of all finite subsets of H .

Let $\text{RT}^{<\mathbb{N}}$ be the statement that for every coloring $f : [\mathbb{N}]^{<\mathbb{N}} \rightarrow 2$ there exists an infinite set H such that $|f([H]^{<\mathbb{N}})| = 1$.

Proposition

$\text{RT}^{<\mathbb{N}}$ is false. (Take f such that $f(s) = 0$ if $|s|$ even and $f(s) = 1$ otherwise.)

Similarly, the generalizations $\text{TS}^{<\mathbb{N}}$, $\text{FS}^{<\mathbb{N}}$ and $\text{RRT}^{<\mathbb{N}}$ are false.
(For $\text{TS}^{<\mathbb{N}}$, take $f(s) = |s|$)

Definition (exactly ω -largeness)

A finite set $s \subseteq \mathbb{N}$ is *exactly ω -large* if $|s| = \min s + 1$.

We write $[H]^{!\omega}$ for the set of all the exactly ω -large subsets of H .

Definition

Let $\text{RT}^{!\omega}$ be the statement: “for every coloring $f : [\mathbb{N}]^{!\omega} \rightarrow 2$ there exists an infinite set H such that $|f([H]^{!\omega})| = 1$.”

Define $\text{TS}^{!\omega}$, $\text{FS}^{!\omega}$ and $\text{RRT}^{!\omega}$ similarly.

Clopen Ramsey theorem

We write $[H]^{\mathbb{N}}$ for the set of all the infinite subsets of H .

- For every infinite set $X = \{x_0, \dots\}$, there exists a unique prefix $\sigma \prec X$ that is exactly ω -large: $x_0, \dots, x_{\min X}$.
- Therefore, for every coloring $f : [\mathbb{N}]^{! \omega} \rightarrow 2$, we can consider the coloring $g : [\mathbb{N}]^{\mathbb{N}} \rightarrow 2$ that send an infinite set X to $f(\sigma)$ for σ its exactly ω -large prefix.
- g is a particular instance of the Clopen Ramsey theorem, which is equivalent to ATR_0 (Friedman/McAloon/Simpson)
- $\text{RT}^{! \omega}$ can also be seen as the first step of a hierarchy of statements (due to Nash-Williams) generalizing RT^n , and concerning coloring of barriers (sets of finite sets having the property that no element is included in another).

Theorem (Carlucci/Gjetaj/L./Levy Patey)

*$TS^{!\omega}$ and $FS^{!\omega}$ proves the existence of $\emptyset^{(\omega)}$, the ω -th jump.
Hence, they imply (and are strictly stronger than) ACA_0 .*

Sketch.

It is sufficient to show the statement for $TS^{! \omega}$ as any free set will be thin.

The statement “for every $f : [\mathbb{N}]^n \rightarrow k$, there is an infinite $H \subseteq \mathbb{N}$ such that $|f([H])^n| \neq k$ ” doesn’t have the cone avoidance property when $k < C_n$ (where C_n is the n -th Catalan number) (Cholak/Patey).

We can get a uniformly computable family of computable coloring $f_n : [\mathbb{N}]^n \rightarrow n$ such that every infinite f_n -thin set computes $\emptyset'$.

We can combine all those instances and obtain some coloring

$f : [\mathbb{N}]^{! \omega} \rightarrow \mathbb{N}$ by taking $f(x_0, \dots, x_{x_0}) = f_{x_0}(x_1, \dots, x_{x_0})$.

Any f -thin set $H = \{x_0, \dots\}$ will be f_{x_i} -thin for every i and therefore computes $\emptyset'$.

We can go further than that and computes $\emptyset^{(\omega)}$ by transforming $\emptyset^{(k)}$ -computable colorings $f : [\mathbb{N}]^n \rightarrow \mathbb{N}$ into computable coloring $f : [\mathbb{N}]^{n+k} \rightarrow \mathbb{N}$ using Δ_k approximations. □

Theorem (Carlucci/Gjetaj/L./Levy Patey)

$\text{RRT}^{!w}$ *have the strong cone avoidance property.*

Proof Sketch

Let C be a non-computable set, and let $f : [\mathbb{N}]^{\omega} \rightarrow \mathbb{N}$ be a coloring such that no color appear more than 2 times.

- For every $s, t \in [\mathbb{N}]^{\omega}$, if $s \neq t$ then $s \not\subseteq t$ (if $\min s \geq \min t$, then $|s| \geq |t|$).
- Let $g : [\mathbb{N}]^{\omega} \rightarrow \mathbb{N}$ be the partial f' -computable coloring defined by $g(s) = \min(t \setminus s)$ if there exists some $t \neq s$ such that $f(t) = f(s)$ and $s <_{\text{lex}} t$.
- g satisfy that either $g(s) > \min s$ or $g(s)$ undefined for every s . We say that g is progressive.
- If H is such that for every $s \in [H]^{\omega}$, $g(s) \notin H$, then for every $s, t \in [H]^{\omega}$, $f(s) \neq f(t)$ otherwise we would have $g(s) = \min(t \setminus s) \in H$.
Hence H is an f -rainbow.

g can be seen as blocking some elements that would prevent a set H from being an f -rainbow.

It is an instance of the free set theorem, but the added assumption that $g(s) > \min s$ allow us to have the strong cone avoidance.

- For every $s \subseteq \mathbb{N}$ such that $|s| \leq \min s$ (i.e. not ω -large), s induce a coloring $f(s, \cdot) : [\mathbb{N}]^{\min s - |s| + 1} \rightarrow \mathbb{N}$
- This is a coloring of tuple of a fixed length.
- We can then use the thin set theorem and the free set theorem for tuple of a fixed size.

The set H is then constructed by forcing.

References



Wei Wang.

Some logically weak Ramseyan theorems.

Adv. Math., 261:1–25, 2014.



Lorenzo Carlucci and Konrad Zdanowski.

The strength of Ramsey's theorem for coloring relatively large sets.

J. Symb. Log., 79(1):89–102, 2014.



Peter Cholak and Ludovic Patey.

Thin set theorems and cone avoidance.

Trans. Amer. Math. Soc., 373(4):2743–2773, 2020.